



Distribution of Mammals in Deramakot & Tangkulap Forest Reserves, Sabah, Malaysia

Hiromitsu Samejima
Robert Ong

Kyoto Working Papers on Area Studies No.129
(G-COE Series 127)

January 2012

The papers in the G-COE Working Paper Series are also available on the G-COE website:
(Japanese webpage)
http://www.humanosphere.cseas.kyoto-u.ac.jp/staticpages/index.php/working_papers
(English webpage)
http://www.humanosphere.cseas.kyoto-u.ac.jp/en/staticpages/index.php/working_papers_en

©2012
Center for Southeast Asian Studies
Kyoto University
46 Shimoadachi-cho,
Yoshida, Sakyo-ku,
Kyoto 606-8501, JAPAN

All rights reserved

ISBN 978-4-906332-01-4

The opinions expressed in this paper are those of the author and do not necessarily reflect the views of the Center for Southeast Asian Studies.

The publication of this working paper is supported by the JSPS Global COE Program (E-04):
In Search of Sustainable Humanosphere in Asia and Africa.

Distribution of Mammals
in Deramakot & Tangkulap Forest Reserves,
Sabah, Malaysia

Hiromitsu Samejima
Robert Ong

Kyoto Working Papers on Area Studies No.129
JSPS Global COE Program Series 127
In Search of Sustainable Humansphere in Asia and Africa

January 2012

Distributions of Mammals in Deramakot & Tangkulap Forest Reserves, Sabah, Malaysia

Hiromitsu Samejima* and Robert Ong†

Introduction

Medium to large mammals of Borneo include various endangered species such as Orangutan and Tembadau and also several important species for the subsistence of local people such as Sambar Deer and Bearded Pig. Thus, these animals are considered as a key element of the forest value and attract international and domestic attentions widely.

Therefore, the distribution of the medium to large mammals in a Forest Management Unit (FMU) is important information for the sustainable management. The information can contribute to select conservation area in the FMU properly, to evaluate performance of current management scheme, and to improve the effectiveness and efficiency of the management. Furthermore, the information also may able to highlight the un-realized economical value of FMU. For example, discovery of high density of orangutan (Ancrenaz et al. 2005) in Malua Forest Reserve, Sabah contributed to launch “the Malua Wildlife Habitat Conservation Bank / Malua BioBank (<http://www.maluabank.com/>)” in 2008 and attract various conservation investments from domestic and oversea entities (Sabah Forestry Department 2008). If a forest management license holder could find a high density area of a charismatic wildlife inside his FMU, he also may be able to develop ecotourism like Danum Valley and Tabin Forest Reserves in that area.

However, there are only a few censuses of mammals over a FMU until now (WWF Malaysia 1982, Ambu 2000). One of the main reasons is the technical difficulty to conduct such a census in a large spatial scale. Therefore, we developed a novel census method: “Random Camera Trapping System”, and conducted in Deramakot and Tangkulap Forest Reserves (827km²), Sandakan, Sabah in 2007-2010. Using this census method, anybody (ex. Non-special stuffs of logging companies) can obtain distribution data of various species economically. In this book, we show the results of our census to exhibit the current situation of the community of medium to large ground-dwelling vertebrates in the FMUs and also the effectiveness of the Random Camera Trapping System as a tool for FMU management.

* Center for Southeast Asian Studies, Kyoto University E-mail:lahang.lejau@gmail.com

† Forest Research Center, Sabah Forestry Department

Deramakot & Tangkulap Forest Reserves

Deramakot and Tangkulap Forest Reserves are continuous forest reserves in Sandakan Division, Sabah, Malaysia at latitude 5°14-30' north and 117°11-35' east (Fig. 1). Both Forest reserves are allocated as Class 2 (Commercial Forest Reserve) and managed as Forest Management Unit (FMU) 19A and part of FMU 17A (with Sungai Pinanga Forest Reserve A). The area is totally 827km² and the altitude is 20-330m above sea level (Fig. 2). Most of the area is covered by lowland mixed dipterocarp forests. However, low-lying area along the Kinabatangan River is covered by freshwater swamp forest (Sabah Forestry Department 2005b, a).




Figure 1. Deramakot (right), Tangkulap (left) and other forest reserves in Sabah. Dark green: total protected forest reserves and national park, Light green: Production forest

Commercial logging was started in 1956 in Deramakot and 1970's in Tangkulap and harvested all the area at least once by conventional logging system. The logging activity was ceased in Deramakot in 1987 and Sustainable Forest Management was started to be conducted by Sabah Forestry Department with cooperation of GTZ (Deutsche Gesellschaft für Technische Zusammenarbeit). The logging in Deramakot was re-started again in 1995, but with Reduced Impact Logging system and have harvested about one fourth of the total area of Deramakot until 2009. On the other hand, conventional logging had been conducted again all over Tangkulap in 1990's and stopped in 2002. Besides, southern parts of Deramakot were burned out during 1983 and 1997 (Sabah Forestry Department 2005b, a).

Sabah Forestry Department drew a Forest Stratum Maps of Deramakot and Tangkulap based on the aerial photos taken in 2001 and SPOT image taken in 2003 respectively (Fig. 3). The forest was classed into four stratum based on densities of large crowns-tree. According to the map, Deramakot has wider range of good and moderate forests (Density of tree DBH ≥ 60 cm is more than 9 tree ha⁻¹) than Tangkulap.

Besides, there are about 11 villages along the Kinabatangan and Milian Rivers south of Deramakot and Tangkulap Forest Reserves. Some of the villages have long history, more than hundreds years. Some of the villagers are used to hunt wildlife as their subsistence foods.




Figure 2. Elevation of Deramakot (right) and Tangkulap (left) Forest Reserves




Figure 3. Forest strata of Deramakot and Tangkulap Forest Reserves

Methods ~Random Camera Trapping System~

We established 29 plots that cover Deramakot and Tangkulap Forest Reserves in a systematic manner (Fig. 4). The plots were about 5 km apart each other. Each plot was a circle with a 1 km diameter. Using the statistical software R 2.10.0 (<http://cran.r-project.org/>), we randomly selected 9–12 setting points to place cameras within each plot. We used an automatic film camera with a passive infrared sensor (Field Note II, Marifu, Iwakuni, Japan) in this study. We first set cameras at three points in each plot and shifted the cameras to the other three points every 3–5 month. We located each setting point using a GPS (GPSmap60CSx, Garmin Ltd., Osoth, KS) and set a camera on a tree about 50 cm above the ground. Each camera faced to open ground and its field-of-view was about 2–7 m². Films and batteries were changed every one to two months.

After we developed the films, we keyed in the data (setting point / date / time / species / no of individuals) of all the photos. We identified the species based on Payne et al. (2005), Myers (2009), and Phillipps and Phillipps (2009). Then we selected data of medium to large mammals (from Moon Rat to Asian Elephant), terrestrial birds and Monitor Lizards. The data of photos captures same species repeatedly in 30 minute were discarded.

For each species, we calculated the numbers of photos taken per unit active camera-days for each plot. The periods when a camera did not work due to full exposure of the film or mechanical failure were excluded from the active periods. We also calculated number of species taken during a 980 camera-days randomly selected in total active camera-days. We iteratively calculated the number 500 times and calculated average number of species for each plots.




Figure 4. 29 plots inside Deramakot and Tangkulap FR. We set 9-12 setting points in each plot.




Figure 5. Setting points of infrared sensor cameras

Results

With increase in active camera days, the number of species taken at each plot has peaked (Fig 6). At each plot, 138-561 photos of 16-27 species were captured in the 980 camera-days (Fig.7). And totally, 7918 photos of 36 species of mammals, 3 species of ground-dwelling birds and one monitor lizard were photographed. In addition, Elying Lemur (*Cynocephalus variegatus*), Slow Loris (*Nycticebus coucang*), Silvered Langur (*Presbytis cristata*), Proboscis Monkey (*Nasalis larvatus*), Bornean Gibbon (*Hylobates muelleri*) and Masked Palm Civet (*Paguma larvata*) are observed by ourselves or recorded by Onoguchi and Matsubayashi (2008) in Deramakot. Thus, 41 species of medium to large mammals were confirmed to inhabit Deramakot in total. They constitute 77.8% of the species recorded in Borneo (Payne and Francis, 2005).




Figure 6. Averaged number of species at each plot




Figure 7. Species composition at each setting area

Species Diversity

The average number of species captured in the 980 camera-days was high at plots in southern and eastern Deramakot (Fig. 8).




Figure 8. Distribution of species richness. Estimated by Inversed Distance Weighted Method with 5 km range

Distribution of individual species

The size of circle shows the amount of mean trapping rates at each plot.


Red list status: Threatened status in IUCN Red list 2011 (IUCN 2011)

Local names: SG: Orang Sungai (Kampung Balat)

Distribution information is based on IUCN (2011)

Moonrat

Echinosorex gymnura




SG: Tikus aputih

Distribution: Peninsular Myanmar, Thailand, Malaysia, Sumatra, and Borneo



Treeshrews




SG: Boasing

Basically arboreal.




Bats



Sunda Pangolin

Manis javanica



Red list status: Endangered

SG: Tengiling


Distribution: Myanmar, Thailand, Laos, Cambodia, Vietnam, Borneo, Sumatra, and Java

The meat and scale are traded in high price and hunting pressure is increasing now.



Western Tarsier

Tarsius bancanus



Red list status: Vulnerable

SG: Lotong lotong


Distribution: Southern Sumatra and Borneo

This species is basically arboreal.



Maroon Leaf Monkey / Red Leaf Monkey

Presbytis rubicunda




Distribution: Borneo endemic

This species is basically arboreal and taken images by chance.



Long-Tailed Macaque / Crab-eating Macaque

Macaca fascicularis



SG: Kara


Distribution: Myanmar, Thailand, Laos, Vietnam
Cambodia, Malaysia, Sumatra, Java, Nusa
Tenggara, and Philippines.

This species is basically arboreal and taken
images by chance.



Southern Pig-Tailed Macaque

Macaca nemestrina



Red list status: Vulnerable


SG: Gebuk

Distribution: Peninsular Thailand, Malaysia, Sumatra and Borneo



Bornean Orangutan

Pongo pygmaeus




Red list status: Endangered


SG: Kagi

Distribution: Endemic to Borneo

This species is known as arboreal, but many adult males walking on forest floor were captured by our cameras.



Squirrels



There are several species of squirrels including Hose-Tailed Squirrel (*Sundasciurus hippurus*), Tufted Ground Squirrel (*Ratufa affinis*), Prevost's Squirrel (*Callosciurus prevostii*), Four Striped Ground Squirrel (*Lariscus hosei*), Giant Squirrel (*Ratufa affinis*), etc.



Hose-Tailed Squirrel



Tufted Ground Squirrel




Prevost's Squirrel




Giant Squirrel

Rats



Common Porcupine

Hystrix brachyura



SG: Oton


Distribution: China, India, Myanmar, Thailand, Laos, Vietnam, Cambodia, Peninsular Malaysia, Sumatra, and Borneo.

Images of two to three individuals walking together are taken sometimes.



Long-Tailed Porcupine

Trichys fasciculata




SG: Oton

Distribution: Peninsular Malaysia, Sumatra, and Borneo



Thick-spined Porcupine

Hystrix crassispinis



SG: Oton


Distribution: Borneo endemic

The distribution pattern in this forest reserves looks like exclusive with common porcupine. Same as common porcupine, images of two to three individuals walking together are taken sometimes.



Sun Bear

Helarctos malayanus



Red list status: Vulnerable


SG: Baruang

Distribution: Myanmar, Thailand, Laos, Vietnam, Cambodia, Peninsular Malaysia, Sumatra, and Borneo



Yellow-throated Marten

Martes flavigula




Distribution: Korea, China, India, Nepal, Myanmar, Thailand, Laos, Vietnam, Cambodia, Peninsular Malaysia, Sumatra, Borneo, and Java

This species is partly arboreal.



Malay weasel

Mustela nudipes




Distribution: Peninsular Thailand, Malaysia, Sumatra, and Borneo

Malay weasel was one of the most elusive species in this area. Only one photo was taken.



Teledu / Malay Badger

Mydaus javanensis




SG: Bilud-tudoh

Distribution: Sumatra, Borneo, and Java

This species is common in Sabah, but rare or does not inhabit in other part of Borneo.



Otters



Red list status: Vulnerable

SG: Rongon


Most of images look like Oriental small-clawed otter (*Aonyx cinerea*). However, Wilting et al. (2010) also reported Smooth-coated otter (*Lutrogale perspicillata*) and Hairy-nosed otter (*Lutra sumatrana*) in this area.

Otters generally inhabit along river.



Malay Civet / Tangalung

Viverra zibellina




SG: Tingalung

Distribution: Peninsular Malaysia, Sumatra, Borneo, Sulawesi and Philippine.



Otter-Civet

Cynogale bennettii



Red list status: Endangered

Distribution: Peninsular Thailand and Malaysia, Sumatra, and Borneo


One of the rarest species in Borneo.

This species is considered to inhabit along river.



Binturong / Bearcat

Arctictis binturong



Red list status: Vulnerable


Distribution: India, Myanmar, Thailand, Lao, Vietnam, Cambodia, Peninsular Malaysia, Sumatra, Borneo, and Java

This species is partly arboreal.



Small-toothed Palm Civet

Arctogalidia trivirgata




Distribution: Northern India, Myanmar, Thailand, Lao, Vietnam, Cambodia, Peninsular Malaysia, Sumatra, and Borneo.

This species is partly arboreal.
Only one photo was taken.



Common Palm Civet

Paradoxurus hermaphroditus



SG: Tingalung

Distribution: Sri Lanka, India, Bangladesh, Myanmar, Thailand, Lao, Vietnam, Cambodia, southern China, Peninsular Malaysia, Sumatra, Borneo, Java, and Philippines.


This species is frequently observed along the logging roads, but was not frequently captured by our cameras inside forest.

This species is partly arboreal.



Banded Civet

Hemigalus derbyanus



Red list status: Vulnerable


SG: Tingalung

Distribution: Peninsular Myanmar, Thailand and Malaysia, Sumatra, and Borneo.



Banded Linsang

Prionodon linsang




Distribution: Peninsular Myanmar, Thailand and Malaysia, Sumatra, Borneo, and Java

This species is partly arboreal.



Collared Mongoose

Herpestes semitorquatus



Red list status: Deficient


Distribution: Endemic to Borneo

Some pictures of this species were very difficult to distinguish from Short-tailed Mongoose.



Short-Tailed Mongoose

Herpestes brachyurus




Distribution: Peninsular Malaysia, Sumatra, and Borneo

Some pictures of this species were very difficult to distinguish from Collared Mongoose.



Sunda Clouded Leopard

Neofelis diardi



Red list status: Vulnerable

SG: Harimau


Distribution: Sumatra and Borneo

This species is the most frequently photographed felid in Deramakot and Tangkulap



Marbled Cat

Pardofelis marmorata




Red list status: Vulnerable

Distribution: Himalayas, Myanmar, Thailand, southern China, Lao, Vietnam, Cambodia, Peninsular Malaysia, Sumatra and, Borneo



Flat-Headed Cat

Prionailurus planiceps



Red list status: Endangered


SG: Kucing hutan

Distribution: Peninsular Malaysia, Sumatra, and Borneo



Leopard Cat

Prionailurus bengalensis



SG: Tusing gauton


Distribution: Northern India, Myanmar, Laos, Cambodia, Vietnam, China, Korea, Taiwan, Peninsular Malaysia, Sumatra, Borneo, Java, Bali, and Palawan

This species is frequently observed along the logging roads, but was not frequently captured by our cameras inside forest.



Bay Cat

Pardofelis badia



Red list status: Endangered


Distribution: Endemic to Borneo

One of the rarest mammals in Borneo



Asian Elephant

Elephas maximus



Red list status: Endangered

SG: Maman / Nenek


Distribution: India, Sri Lanka, Myanmar, Thailand, Lao, Southern China, Vietnam, Cambodia, Peninsular Malaysia, Sumatra. and Borneo

Many cameras were destroyed by elephants.



Bearded Pig

Sus barbatus



Red list status: Vulnerable

SG: Bakas


Distribution: Peninsular Malaysia, Sumatra, and Borneo

During mast fruiting season such as in 2007 and 2010, their density increased obviously and large herds including more than 20 piglets were observed.



Lesser Mousedeer & Greater Mousedeer

Tragulus javanicus & *Tragulus napu*



SG: Lesser: Palanuk, Greater: Palanuk watang

Distribution:

Lesser: Thailand, Laos, Cambodia, Vietnam, Peninsular Malaysia, Sumatra, and Borneo

Greater: Peninsular Myanmar, Thailand and Malaysia, Sumatra, and Borneo

Some pictures of these two species were very difficult to distinguish.




Lesser Mousedeer



Greater Mousedeer

Bornean Yellow Muntjac

Muntiacus atherodes



SG: Parukak


Distribution: Endemic to Borneo

The skin colors vary from yellow to brown. Some people reported inhabitation of Bornean Red Muntjac (*Muntiacus muntjac*) in Deramakot. However, we could not get any obvious picture of that species which is characterized by the two-spikes-antlers.



Sambar Deer

Rusa unicolor



Red list status: Vulnerable


SG: Payaw

Distribution: Sri Lanka, India, southern China, Taiwan, Myanmar, Thailand, Laos, Vietnam, Cambodia, Sumatra, and Borneo



Tembadau / Banteng

Bos javanicus



Red list status: Endangered

SG: Kalasiw


Distribution: Myanmar, Thailand, Cambodia, Borneo, and Java

Some informants said, domestic cattle (*Bos indicus*) were once bred at a logging camp near Kg Balat. And some of the cattle were escaped into the forest when the camp was closed in 1980's. Some part of wild cows in Deramakot & Tangkulap may be their offspring or hybrid with wild Tembadau.



Chestnut Necklaced Partridge

Arborophila charltonii graydoni




Red list status: Near threatened

Distribution: Peninsular Thailand and Malaysia, Sumatra, and Borneo (only in Sabah)



Crested Fireback

Lophura ignita nobilis




Red list status: Near threatened

Distribution: Peninsular Thailand and Malaysia, Sumatra, and Borneo



Great Argus

Argusianus argus grayi



Red list status: Near threatened

SG: Manuk gautun

Distribution: Peninsular Thailand and Malaysia, Sumatra, and Borneo



Other Bird species


Many other bird species were also photographed. The identified species are Red-breasted Partridge (*Arborophila hyperythra*), Crested Partridge (*Rollulus rouloul*), Crested Wood Partridges (*Rollulus rouloul*), Red-legged Crake (*Rallina fasciata*), Band-bellied Crake (*Porzana paykullii*), Storm's Stork (*Ciconia stormi*), Yellow Bittern (*Ixobrychus sinensis*), Blyth's Hawk Eagle (*Spizaetus alboniger*), Emerald Dove (*Chalcophaps indica*), Pink-necked Green Pigeon (*Treron vernans*), Rufous-collared Kingfisher (*Actenoides concretus*), White-crowned Hornbill (*Berenicornis comatus*), Blue-headed Pitta (*Pitta baudii*), Black-and-crimson Pitta (*Pitta ussheri*), Giant Pitta (*Pitta granatina*), Hooded Pitta (*Pitta sordida*), Blue-banded Pitta (*Pitta arquata*), Banded Pitta (*Pitta guajana*) Garnet Pitta (*Pitta granatina*), Bornean Ground-cuckoo (*Carpococcyx radiates*), Black-capped Babbler (*Pellorneum capistratum*), White-crowned Shama (*Copsychus stricklandii*), Bornean Ground-babbler (*Ptilochila leucogrammica*).



Black rough-neck Monitor Lizard

& Common Water Monitor Lizard

Varanus rudicollis & *V. salvator*



SG: Mandatan



Black rough-neck Monitor Lizard



Common Water Monitor Lizard

Other reptile



Acknowledgments

We deeply appreciate to Datuk Sam Mannan, the Director of Sabah Forestry Department and Prof. Kanehiro Kitayama (Kyoto University) for their kindly support of the project. We appreciate to all the staffs of Deramakot Forest Reserve especially Mr. Subari Suparlan, Peter Lagan, Azny Ahmad, Edward Lapina, Edward Thomas, Andaus Bagoi, Sah Joo Achang, Isran Mahran and Robin Dulasag who helped our field work and informed the management system. We also appreciate many staffs of Forestry Research Center, especially Mr. Julsin Sikui and Mohamad Jumri and also Mr. Dimeh Koyopo of Sabah Forestry Department. We thanks to Dr. Hisashi Matsubayashi (Universiti Malaysia Sabah), Yoshihiro Nakashima (Kyoto University), Ms. Miyabi Nakabayashi (Kyoto University) and Mr. Go Onoguchi for their technical help, Mr. Muniandy@Andy Marcus, Ronnie Marcus, Jonik Ajis, Ronald Samion, Jubilee bin George, Sandra Lakim, Azrie Bernad, Rawinder Ajon and Sidik Ajis for their hard field work and Suhayli bin Ardeh for information of local names.

This work was partly supported by the Environment Research and Technology Development Fund (F-071 and D-1006) of the Ministry of the Environment, Japan and Grant-in-Aid for Scientific Research (S) 810104300001 of Japan Society for the Promotion of Science. Publication of this working paper is supported by JSPS Global COE Program “In Search of Sustainable Humanosphere in Asia and Africa” in Center for Southeast Asian Studies, Kyoto University.

Local assistants for the hard filed works: Andy Marcus, Ronnie Marcus, Ronald Samion , Jubilee bin George, Sandra Lakim, Jonik Ajis, Azrie, Ewin and Sidik Ajis (left to right)



References

- Ambu, L. N. 2000. Survey of part of the Sapulut Forest Reserve and Ulu S. Milian Forest Reserve (FMU).in D. Aloysius, R. Goddos, and F. Johniu, editors. 4th Seminar on Tropical Ecosystem Research in Sabah "Research Contributions towards Optimisation of Researchs in Tropical Ecosystems". Institute for Development Studies (Sabah), Kota Kinabalu.
- Ancrenaz, M., O. Gimenez, L. Ambu, K. Ancrenaz, P. Andau, B. Goossens, J. Payne, A. Sawang, A. Tuuga, and I. Lackman-Ancrenaz. 2005. Aerial surveys give new estimates for orangutans in Sabah, Malaysia. PLoS Biology 3:1-8.
- Myers, S. 2009. *A field guide to the birds of Borneo*. Talisman Publishing, Singapore.
- Onoguchi, G. and H. Matsubayashi. 2008. Comparative study on mammalian fauna in different harvesting intensities with reduced-impact and conventional logging in Sabah, Malaysia. Pages 133-140 in Research Institute for Humanity and Nature Project_2-2, editor. Sustainability and biodiversity assessment on forest utilization options.
- Payne, J., C. M. Francis, and K. Phillips. 2005. *A Field Guide to the Mammals of Borneo*. The Sabah Society, Kota Kinabalu.
- Phillipps, Q. and K. Phillipps. 2009. *Phillipps' Field Guide to the Birds of Borneo*. Beaufoy Books, Oxford.
- Sabah Forestry Department. 2005a. *Environmental Impact Assessment for Tangkulap-Pinangah Forest Development Forest Management Unit 17 Part A (FMU17 Part A)*.
- Sabah Forestry Department. 2005b. *Forest Management Plan 2 (1st January 2005 - 31th December 2014) Deramakot Forest Reserve Forest Management Unit No. 19*.
- Sabah Forestry Department. 2008. *Annual Report 2007*.
- Wilting, A., H. Samejima, and A. Mohamed. 2010. Diversity of Bornean viverrids and other small carnivores in Deramakot Forest Reserve, Sabah. *Small Carnivore Conservation* 42:10-13.
- WWF Malaysia. 1982. *A Faunal Survey of Sabah*.